# Biotinylated Human IL-22R alpha 1&IL-10R beta Protein





Description	
Source	Recombinant Biotinylated Human IL-22R alpha 1&IL-10R beta Protein is expressed from HEK293 with hFc tag and Avi Tag at the C-Terminus.
	It contains His16-Thr228 (IL-22R alpha 1) and Met20-Ser220 (IL-10R beta).
Accession	Q8N6P7(IL-22R alpha 1)&Q08334(IL-10R beta)
Molecular Weight	The protein has a predicted MW of 52.07 kDa (IL-22R alpha 1) and 48.85 kDa (IL-10R beta). Due to glycosylation, the protein migrates to 68-75 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
	> 95% as determined by HPLC
Formulation and Storage	

#### Formulation and Storage


Formulation	Lyophilized from 0.22μm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 24 months as supplied from date of receipt80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

# **Background**

IL-22R $\alpha$ 1&IL-10R $\beta$  are receptors for IL-22 and IL-10, mediate the activation of JAK-STAT, MAPK and other signaling pathways, and participate in the process of many inflammatory diseases. IL-22 R $\alpha$ 1 contains cytoplasmic motifs for interactions with signal transduction molecules, but formation of ternary complexes with IL10 R $\beta$  or IL20 R $\beta$  and the respective ligands is required for signal transduction[1].


## **Assay Data**

#### **Bis-Tris PAGE**



Biotinylated Human IL-22R alpha 1&IL-10R beta on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

#### **SEC-HPLC**



The purity of Biotinylated Human IL-22R alpha 1&IL-10R beta is greater than 95% as determined by SEC-HPLC.




## **Assay Data**

#### **ELISA Data**

# Biotinylated Human IL-22R alpha 1&IL-10R beta, hFc Tag ELISA

 $0.5 \mu g$  Human IL-22, His Tag Per Well



Log Biotinylated Human IL-22R alpha 1&IL-10R beta, hFc Tag Conc.( $\mu g/ml$ )

Immobilized Human IL-22, His Tag at  $5\mu g/ml$  (100 $\mu$ I/well) on the plate. Dose response curve for Biotinylated Human IL-22R alpha1&IL-10R beta, hFc Tag with the EC50 of  $1.36\mu g/ml$  determined by ELISA.