# Mouse IL-13Ra2 Protein

#### Cat. No. ILR-MM1R2



Description	
Source	Recombinant Mouse IL-13Ra2 Protein is expressed from HEK293 with His tag at the C-terminus.
	It contains Leu22-Lys334.
Accession	O88786-1
Molecular Weight	The protein has a predicted MW of 37.55 kDa. Due to glycosylation, the protein migrates to 45-60 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
	> 95% as determined by HPLC

#### Formulation and Storage


Formulation	Lyophilized from 0.22 µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt80°C for 3-6 months after reconstitution. 2-8°C for 2-7 days after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

# **Background**

Two type 1 membrane proteins belonging to the hemopoietin receptor family have been cloned and shown to bind IL-13 with differing affinities. The lower affinity IL-13 binding protein, previously designated IL-13 R alpha, IL-13 R alpha ' or NR4, is now referred to as IL-13 R alpha 1. The high-affinity IL-13 binding protein, previously also designated IL-13 R or IL-13 R alpha ', is now referred to as IL-13 R alpha 2.

# **Assay Data**

#### **Bis-Tris PAGE**




Mouse IL-13Ra2 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

**SEC-HPLC** 

# KAGTUS


# **Assay Data**



The purity of Mouse IL-13Ra2 is greater than 95% as determined by SEC-HPLC.

#### **ELISA Data**

Mouse IL-13R2, His Tag ELISA 0.1µg Mouse IL-13R2, His Tag Per Well



Immobilized Mouse IL-13R2, His Tag at 1µg/ml (100µl/well) on the plate. Dose response curve for Human IL-13, hFc Tag with the EC50 of 6.6ng/ml determined by ELISA.