#### Cynomolgus Integrin alpha 4 beta 7 (ITGA4&ITGB7) Heterodimer Protein

Cat. No. ITG-CM147



Weightmigrates to 73-80 kDa, 110-120 kDa and 130-150 kDa based on Bis-Tris PAGE result.EndotoxinLess than 0.1 EU per μg by the LAL method.	Description	
Accession XP_005573683.3(ITGA4)&XP_005571028.3(ITGB7)  Molecular Weight The protein has a predicted MW of 110.87 kDa (ITGA4) and 81.09 kDa (ITGB7). Due to glycosylation, the protein migrates to 73-80 kDa, 110-120 kDa and 130-150 kDa based on Bis-Tris PAGE result.  Endotoxin Less than 0.1 EU per μg by the LAL method.	Source	· · · · · · · · · · · · · · · · · · ·
Molecular Weight  The protein has a predicted MW of 110.87 kDa (ITGA4) and 81.09 kDa (ITGB7). Due to glycosylation, the protein migrates to 73-80 kDa, 110-120 kDa and 130-150 kDa based on Bis-Tris PAGE result.  Endotoxin  Less than 0.1 EU per μg by the LAL method.		It contains Tyr34-Thr977 (ITGA4) acidic tail and Glu20-His723 (ITGB7) basic tail.
Weightmigrates to 73-80 kDa, 110-120 kDa and 130-150 kDa based on Bis-Tris PAGE result.EndotoxinLess than 0.1 EU per μg by the LAL method.	Accession	XP_005573683.3(ITGA4)&XP_005571028.3(ITGB7)
2		The protein has a predicted MW of 110.87 kDa (ITGA4) and 81.09 kDa (ITGB7). Due to glycosylation, the protein migrates to 73-80 kDa, 110-120 kDa and 130-150 kDa based on Bis-Tris PAGE result.
Purity > 95% as determined by Ris-Tris PAGE	Endotoxin	Less than 0.1 EU per μg by the LAL method.
2 00% do determined by bio-Thor Ace	Purity	> 95% as determined by Bis-Tris PAGE

#### Formulation and Storage

Formulation Supplied as 0.22 μm filtered solution in PBS (pH 7.4).

Storage Valid for 12 months from date of receipt when stored at -80°C. Recommend to aliquot the protein into smaller


quantities for optimal storage. Please minimize freeze-thaw cycles.

#### **Background**

Integrins alpha-4/beta-1 (VLA-4) and alpha-4/beta-7 are receptors for fibronectin. They recognize one or more domains within the alternatively spliced CS-1 and CS-5 regions of fibronectin. They are also receptors for VCAM1. Integrin alpha-4/beta-1 recognizes the sequence Q-I-D-S in VCAM1. Integrin alpha-4/beta-7 is also a receptor for MADCAM1. It recognizes the sequence L-D-T in MADCAM1.

#### **Assay Data**

#### **Bis-Tris PAGE**



Cynomolgus ITGA4&ITGB7 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

#### **ELISA Data**

# **Cynomolgus ITGA4&ITGB7, His Tag ELISA** 0.2μg Cynomolgus ITGA4&ITGB7, His Tag Per Well

3.5 3.0 2.5 0.5 0.5 0.0 -4 -3 -2 -1 0 Log Anti-ITGA4 Antibody, hFc Tag Conc.(µg/ml)

Immobilized Cynomolgus ITGA4&ITGB7, His Tag at  $2\mu g/ml$  (100 $\mu l/well$ ) on the plate. Dose response curve for Anti-ITGA4 Antibody, hFc Tag with the EC50 of 4.2ng/ml determined by ELISA.


### Cynomolgus Integrin alpha 4 beta 7 (ITGA4&ITGB7) Heterodimer Protein

Cat. No. ITG-CM147



## **Assay Data**

#### **BLI Data**



Loaded Anti-ITGA4 Antibody, hFc Tag on ProA-Biosensor can bind Cynomolgus ITGA4&ITGB7, His Tag with an affinity constant of 17.80 nM as determined in BLI assay (Gator® Prime).