

Cynomolgus Integrin alpha 2 beta 1 (ITGA2&ITGB1) Heterodimer Protein

Cat. No. ITG-CM1AB

Description

Source	Recombinant Cynomolgus Integrin alpha 2 beta 1 (ITGA2&ITGB1) Heterodimer Protein is expressed from HEK293 with His tag at the C-terminus. It contains Tyr30-Pro1130 (ITGA2) acidic tail and Gln161-Asp868 (ITGB1) basic tail.
Accession	G8F2Z5(ITGA2)&A0A7N9D0D7(ITGB1)
Molecular Weight	The protein has a predicted MW of 126.69 kDa (ITGA2) and 83.16 kDa (ITGB1). Due to glycosylation, the protein migrates to 130-170 kDa (ITGA2) and 100-130 kDa (ITGB1) based on Tris-Bis PAGE result.
Endotoxin	Less than 1EU per μ g by the LAL method.
Purity	> 95% as determined by Tris-Bis PAGE > 95% as determined by HPLC

Formulation and Storage


Formulation	Lyophilized from 0.22 μ m filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 μ g/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3-6 months after reconstitution. 2-8°C for 2-7 days after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

The α 2 β 1 integrin, also known as VLA-2, GPIa-IIa, CD49b, was first identified as an extracellular matrix receptor for collagens and/or laminins. It is now recognized that the α 2 β 1 integrin serves as a receptor for many matrix and nonmatrix molecules. It plays a critical role in platelet function and homeostasis.


Assay Data

Tris-Bis PAGE



Cynomolgus ITGA2&ITGB1 on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC



The purity of Cynomolgus ITGA2&ITGB1 is greater than 95% as determined by SEC-HPLC.