
Human Integrin alpha 5 beta 1 (ITGA5&ITGB1) Heterodimer Protein, Ultra Lo w Endotoxin



| Cat. No. | ITG-HM451- | |
|-------------------------|------------|---|
| Description | | |
| Source | | Recombinant Human Integrin alpha 5 beta 1 (ITGA5&ITGB1) Heterodimer Protein is expressed from HEK293 with His tag and Avi tag at the C-Terminus of ITGA5. |
| | | It contains Phe42-Tyr995(ITGA5) acidic tail and Gln21-Asp728(ITGB1) basic tail. |
| Accession | | P08648(ITGA5)&P05556-1(ITGB1) |
| Molecular Weight | | The protein has a predicted MW of 111.8 kDa (ITGA5) & 83.2 kDa (ITGB1). Due to glycosylation, the protein migrates to 100-140 kDa based on Bis-Tris PAGE result. |
| Endotoxin | | Less than 0.01 EU per μg by the LAL method. |
| Purity | | > 95% as determined by Bis-Tris PAGE |
| Fully | | > 95% as determined by HPLC |
| Formulation and Storage | | |
| Formulatic | n | Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization. |
| Reconstitu | ition | Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions. |
| Storage | | -20 to -80°C for 24 months as supplied from date of receipt80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles. |
| Background | | |
| | | Integrin alpha 5/ beta 1, also known as VLA-5, is a widely expressed non-covalent heterodimer of a 160 kDa alpha 5 and a 130 kDa beta 1 Integrin subunit.Alpha 5/ beta 1 is upregulated on tumor vasculature and promotes angiogenesis. |


Assay Data

Bis-Tris PAGE



Human ITGA5&ITGB1 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

Human Integrin alpha 5 beta 1 (ITGA5&ITGB1) Heterodimer Protein, Ultra Lo w Endotoxin




The purity of Human ITGA5&ITGB1 is greater than 95% as determined by SEC-HPLC.

KVCJUS

ELISA Data

Human ITGA5&ITGB1, His Tag ELISA





Immobilized Human ITGA5&ITGB1, His Tag at 1μ g/ml (100 μ l/well) on the plate. Dose response curve for Anti-ITGA5&ITGB1 Antibody, hFc Tag with the EC50 of 16.6ng/ml determined by ELISA (QC Test).