

Biotinylated Human KIR2DL2 Protein

Cat. No. KIR-HM4L2B

Description

Source	Recombinant Biotinylated Human KIR2DL2 Protein is expressed from HEK293 with His tag and Avi tag at the C-Terminus. It contains His22-His245.
Accession	P43627
Molecular Weight	The protein has a predicted MW of 27.4 kDa. Due to glycosylation, the protein migrates to 45-60 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE > 90% as determined by HPLC

Formulation and Storage


Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

KIR2DL2 (2DL2, formerly NKAT6, designated CD158b) is a 348 amino acid (aa) type I transmembrane glycoprotein that belongs to the human killer cell Iglike receptor (KIR) family. KIRs are expressed on human CD56dim NK cells and T cell subsets, and regulate effector functions in the innate immune system. KIR2DL2 is receptor on natural killer (NK) cells for HLA-Cw1, 3, 7, and 8 allotypes. Inhibits the activity of NK cells thus preventing cell lysis.


Assay Data

Bis-Tris PAGE



Biotinylated Human KIR2DL2 on Bis-Tris PAGE under reduced conditions. The purity is greater than 95%.

SEC-HPLC



The purity of Biotinylated Human KIR2DL2 is greater than 90% as determined by SEC-HPLC.