

Human Kallikrein 3/PSA Protein (active form)



Cat. No. KLK-HM003

Description	
Source	Recombinant Human Kallikrein 3/PSA Protein (active form) is expressed from HEK293 without tag. It contains Ile25-Pro261.
Accession	P07288-1
Molecular Weight	The protein has a predicted MW of 26.08 kDa. Due to glycosylation, the protein migrates to 28-38 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1 EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC

Formulation and Storage	
Formulation	Lyophilized from 0.22 µm filtered solution in 50mM Tris, 150mM NaCl, 0.05% Brij-35 (pH 7.5). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

Kallikrein-related peptidase 3 (KLK3), also known as prostate-specific antigen (PSA), is the most useful biomarker for prostate cancer (PCa). KLK3 is suggested to play a role in regulating cancer growth through anti-angiogenic activity in vivo and in vitro.


Assay Data

Bis-Tris PAGE



Human Kallikrein 3 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC



The purity of Human Kallikrein 3 is greater than 95% as determined by SEC-HPLC.

Human Kallikrein 3/PSA Protein (active form)

Cat. No. KLK-HM003




Assay Data

ELISA Data

Human Kallikrein 3, No Tag ELISA

0.05µg Human Kallikrein 3, No Tag Per Well



Immobilized Human Kallikrein 3, No Tag at 0.5µg/ml (100µl/well) on the plate. Dose response curve for Anti-Kallikrein 3 Antibody, hFc Tag with the EC50 of 7.3ng/ml determined by ELISA.

Bioactivity Data

Measured by its ability to cleave the colorimetric peptide substrate, Succinyl-Arg-Pro-Tyr-p-Nitroanilide (Suc-RPY-pNA). The specific activity is >500 pmol/min/µg.