
## Biotinylated Human Kallikrein 7/KLK7 Protein (pro form)

# ᠺ᠕ᢏᠴ᠊᠐᠍᠋ᠶ


Cat. No.	KLK-HM417	лв ПОЛССТОЗ
Description		
Source		Recombinant Biotinylated Human Kallikrein 7/KLK7 Protein (pro form) is expressed from HEK293 with His tag and Avi Tag at the C-terminus. The protein needs to be activated by Thermolysin for an activated form.
		It contains Glu23-Arg253.
Accession		P49862
Molecular Weight		The protein has a predicted MW of 28.11 kDa. Due to glycosylation, the protein migrates to 30-40 kDa based on Bis-Tris PAGE result.
Endotoxin		Less than 1 EU per μg by the LAL method.
Durit (		> 95% as determined by Bis-Tris PAGE
Purity		> 95% as determined by HPLC
Formulation and Storage		
Formulatio	n	Supplied as 0.22 µm filtered solution in 20mM HEPES, 150mM NaCL (pH 7.5).
Storage		Valid for 12 months from date of receipt when stored at -80°C. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.
Background		
		Kallikrein 7 (KLK7) is a secreted serine protease with chymotrypsic protease activity. Abnormally high expression of KLK7 is closely related to the occurrence and development of various types of cancer. Therefore, KLK7 has been identified as a potential target for cancer drug development design in recent years. KLK7 mediates various biological and pathological processes in tumorigenesis, including cell proliferation, migration, invasion, angiogenesis, and cell metabolism, by hydrolyzing a series of substrates such as membrane proteins, extracellular matrix proteins, and cytokines.

Assay Data

### **Bis-Tris PAGE**



#### SEC-HPLC



Biotinylated Human Kallikrein 7 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

The purity of Biotinylated Human Kallikrein 7 is greater than 95% as determined by SEC-HPLC.

## Biotinylated Human Kallikrein 7/KLK7 Protein (pro form)

Cat. No. KLK-HM417B



**Bioactivity Data** 

Measured by its ability to cleave the fluorogenic peptide substrate, Mca-RPKPVE-Nval-WRK(Dnp)-NH2. The specific activity is >150 pmol/min/µg.

