### Mouse KLRG1 Protein

#### Cat. No. KLR-MM2G1



Description	
Source	Recombinant Mouse KLRG1 Protein is expressed from HEK293 with hFc tag at the N-Terminus.
	It contains Gln57-Tyr188.
Accession	O88713
Molecular Weight	The protein has a predicted MW of 42.3 kDa. Due to glycosylation, the protein migrates to 50-65 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
	> 95% as determined by HPLC

### Formulation and Storage

minimize freeze-thaw cycles.


Formulation	Lyophilized from 0.22 µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt80°C for 3-6 months after reconstitution.2-8°C for 2-7 days after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please

## **Background**

Immune homeostasis requires the tight, tissue-specific control of the different CD4 Foxp3 regulatory T (Treg) cell populations. The cadherin-binding inhibitory receptor killer cell lectin-like receptor G1 (KLRG1) is expressed by a subpopulation of Treg cells with GATA3 effector phenotype.Lack of KLRG1 on Treg cells conferred on them a competitive advantage in the gut, but not in lymphoid organs. Hence, although absence of KLRG1 is not enough to increase intestinal Treg cells in KLRG1 knockout mice, KLRG1 ligation reduces T-cell receptor signals and the competitive fitness of individual Treg cells in the intestine.

#### **Assay Data**

#### **Bis-Tris PAGE**




Mouse KLRG1 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

**SEC-HPLC** 



# **Assay Data**



The purity of Mouse KLRG1 is greater than 95% as determined by SEC-HPLC.