Cynomolgus Kremen-2 Protein

Cat. No. KRE-CM102



Description	
Source	Recombinant Cynomolgus Kremen-2 Protein is expressed from HEK293 with His tag at the C-Terminus.
	It contains Gly26-Ala364.
Accession	XP_005591068.2
Molecular Weight	The protein has a predicted MW of 37.04 kDa. Due to glycosylation, the protein migrates to 50-65 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
	> 95% as determined by HPLC

Formulation and Storage


· · · · · · · · · · · · · · · · · · ·	
Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

Kremen2 (Krm2) plays an important role in embryonic development, bone formation, and tumorigenesis as a crucial regulator of classical Wnt/β-catenin signaling pathway. Compared to para-cancerous tissues, Krm2 was significantly up-regulated in gastric cancer tissues and was positively correlated with the pathological grade of gastric cancer patients. Krm2 can be a potent candidate for designing of targeted therapy.


Assay Data

Bis-Tris PAGE



Cynomolgus Kremen-2 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC




The purity of Cynomolgus Kremen-2 is greater than 95% as determined by SEC-HPLC.

KAGTUS

Assay Data

ELISA Data

Cynomolgus Kremen-2, His Tag ELISA 0.05µg Cynomolgus Kremen-2, His Tag Per Well



Immobilized Cynomolgus Kremen-2, His Tag at $0.5\mu g/ml$ (100 μ l/well) on the plate. Dose response curve for Anti-Kremen-2 Antibody, hFc Tag with the EC50 of 4.8ng/ml determined by ELISA.