# Biotinylated Human LAIR2/CD306 Protein





Description	
Source	Recombinant Biotinylated Human LAIR2/CD306 Protein is expressed from HEK293 with His tag and Avi tag at the C-Terminus.
	It contains Gln22-Pro152.
Accession	Q6ISS4-1
Molecular Weight	The protein has a predicted MW of 17 kDa. Due to glycosylation, the protein migrates to 20-25 kDa based on Bis- Tris PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
	> 95% as determined by HPLC

#### Formulation and Storage


Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

# Background

The leukocyte-associated immunoglobulin-like receptor 2 (LAIR2) shows increased expression during pregnancy and its reduced expression is associated with pregnancy complications. Although LAIR2 was expressed in the human placenta villus and decidua in the first trimester of pregnancy, it was not expressed in mouse tissues. Knockdown of LAIR2 markedly improved cell viability and inhibited the invasive ability of HTR8/SVneo cells.


## **Assay Data**

#### **Bis-Tris PAGE**



Biotinylated Huma LAIR2 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

#### **SEC-HPLC**



The purity of Biotinylated Huma LAIR2 is greater than 95% as determined by SEC-HPLC.


# Biotinylated Human LAIR2/CD306 Protein

Cat. No. LAI-HM4R2B

# KAGTUS

# **Assay Data**

## **SPR Data**



Biotinylated Human LAIR2, His-Avi Tag immobilized on CM5 Chip can bind Native Human Collagen I, No Tag with an affinity constant of 1.80 nM as determined in SPR assay (Biacore T200).