# Biotinylated Human LAP (TGF beta 1) Protein





| Description         |  |
|---------------------|--|
| Source              | Recombinant Biotinylated Human LAP (TGF beta 1) Protein is expressed from HEK293 with His tag and Avi tag at the N-Terminus.       |
|                     | It contains Leu30-Arg278(C33S).  |
| Accession           | P01137   |
| Molecular<br>Weight | The protein has a predicted MW of 31.4 kDa. Due to glycosylation, the protein migrates to 45-50 kDa based on Bis-Tris PAGE result. |
| Endotoxin           | Less than 1EU per μg by the LAL method.  |
| Purity              | > 95% as determined by Bis-Tris PAGE   |

# Formulation and Storage

Formulation Supplied as 0.22µm filtered solution in PBS (pH 7.4).

Storage Valid for 12 months from date of receipt when stored at -80°C. Recommend to aliquot the protein into smaller


quantities for optimal storage. Please minimize freeze-thaw cycles.

# **Background**

Latent TGF beta 1 cDNA encodes a 390 amino acid precursor that contains a 29 aa signal peptide and a 361 aa proprotein. A furinlike convertase processes the proprotein to generate an Nterminal 249 aa latencyassociated peptide (LAP) and a Cterminal 112 aa mature TGF beta 1. Disulfidelinked homodimers of LAP and TGF beta 1 remain noncovalently associated after secretion, forming the small latent TGF beta 1 complex.

### **Assay Data**

#### **Bis-Tris PAGE**




Biotinylated Human LAP (TGF beta 1) on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

### **ELISA Data**

### Biotinylated Human LAP (TGF beta 1), His Tag ELISA

0.2μg Human ITGAV&ITGB6, His Tag Per Well



Log Biotinylated Human LAP (TGF beta 1), His Tag Conc.(μg/ml)

Immobilized Human ITGAV&ITGB6, His Tag at 2 $\mu$ g/ml (100 $\mu$ l/well) on the plate. Dose response curve for Biotinylated Human LAP (TGF beta 1) , His Tag with the EC50 of 0.11 $\mu$ g/ml determined by ELISA (QC Test).