

Canine LIF R/CD118 Protein, Ultra Low Endotoxin



Cat. No. LIF-DM10R-UL


Description	
Source	Recombinant Canine LIF R/CD118 Protein is expressed from HEK293 with His tag at the C-terminus. It contains Glu45-Ser833.
Accession	AAU43788.1
Molecular Weight	The protein has a predicted MW of 91.21 kDa. Due to glycosylation, the protein migrates to 120-150 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 0.01 EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC

Formulation and Storage	
Formulation	Lyophilized from 0.22 µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background	
The leukemia inhibitory factor receptor (LIF-R) subunit is a component of cell-surface receptor complexes for the multifunctional cytokines, LIF, cardiotrophin-1, ciliary neurotrophic factor, and human oncostatin M. The structure of the human LIF-R gene is similar to that of the mouse gene. The transmembrane receptor is encoded by 19 exons.	

Assay Data


Bis-Tris PAGE



Canine LIF R on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

Assay Data




The purity of Canine LIF R is greater than 95% as determined by SEC-HPLC.

ELISA Data

Canine LIF R, His Tag ELISA

0.1µg Human LIF, No Tag Per Well



Immobilized Human LIF, No Tag at 1µg/ml (100µl/well) on the plate. Dose response curve for Canine LIF R, His Tag with the EC50 of 25.4ng/ml determined by ELISA.