

Human LILRB2/CD85d/ILT4 Protein, Ultra Low Endotoxin

Cat. No. LIL-HM2B2-UL

Description

Source	Recombinant Human LILRB2/CD85d/ILT4 Protein is expressed from HEK293 with hFc tag at the C-terminus. It contains Gln22-Arg457.
Accession	AAL36991.1
Molecular Weight	The protein has a predicted MW of 74.1 kDa. Due to glycosylation, the protein migrates to 80-110 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 0.01 EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC

Formulation and Storage


Formulation	Lyophilized from 0.22µm filtered solution in 20mM Tris, 100mM glycine, 100mM L-arginine, 150mM NaCl (pH 7.5) / PBS, 200mM L-Arginine (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

The immunoglobulin-like transcript (ILT) comprise a family of activating and inhibitory type immunoreceptors whose genes are located in the same locus that encodes killer cell Ig-like receptors (KIR). ILT4, also known as LIR-2 and LILRB2, is a type I transmembrane protein expressed primarily on monocytes and dendritic cells (DC). LILRB2 is a receptor for class I MHC antigens. Recognizes a broad spectrum of HLA-A, HLA-B, HLA-C, HLA-G and HLA-F alleles.


Assay Data

Bis-Tris PAGE



Human LILRB2 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC




The purity of Human LILRB2 is greater than 95% as determined by SEC-HPLC.

Assay Data


ELISA Data

Human LILRB2, hFc Tag ELISA
0.1µg Human LILRB2, hFc Tag Per Well




Immobilized Human LILRB2, hFc Tag at 1µg/ml (100µl/well) on the plate. Dose response curve for Biotinylated Anti-LILRB2 Antibody, hFc Tag with the EC₅₀ of 14.6ng/ml determined by ELISA (QC Test).

SPR Data



Human LILRB2, hFc tag captured on Protein A Chip can bind Human HLA-G Tetramer, His Tag with an affinity constant of 9.20nM as determined in SPR assay (Biacore T200).

SPR Data



Human LILRB2, hFc Tag captured on Protein A chip, can bind Human ANGPTL2, His Tag with an affinity constant of 0.14µM as determined in a SPR assay (Biacore T200).