Human LRP-5 Protein

Cat. No. LRP-HM305



Description	
Source	Recombinant Human LRP-5 Protein is expressed from HEK293 with mFc (IgG1) tag at the C-Terminus.
	It contains Glu644-Gln1263.
Accession	O75197-1
Molecular Weight	The protein has a predicted MW of 96 kDa. Due to glycosylation, the protein migrates to 100-115 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE

Formulation and Storage

Formulation Supplied as 0.22µm filtered solution in PBS (pH 7.4).

Storage Valid for 12 months from date of receipt when stored at -80°C. Recommend to aliquot the protein into smaller


quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

A role for low-density lipoprotein-related receptor 5 (LRP5) in human bone was first established by the identification of genetic alterations that led to dramatic changes in bone mass. Shortly thereafter, mutations that altered the function of the sclerostin (SOST) gene were also associated with altered human bone mass. Subsequent studies of LRP5 and sclerostin have provided important insights into the mechanisms by which these proteins regulate skeletal homeostasis.

Assay Data

Bis-Tris PAGE




Human LRP-5 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

ELISA Data

Human LRP-5, mFc Tag ELISA

 $0.5\mu g$ Human Wnt Surrogate-Fc Fusion, hFc Tag Per Well



Immobilized Human Wnt Surrogate-Fc Fusion, hFc Tag at $5\mu g/ml$ ($100\mu l/well$) on the plate. Dose response curve for Human LRP-5, mFc Tag with the EC50 of $0.73\mu g/ml$ determined by ELISA (QC Test).




Assay Data

ELISA Data

Human LRP-5, mFc Tag ELISA

0.5μg Human DKK1, His Tag Per Well



Immobilized Human DKK1, His Tag at $5\mu g/ml$ (100 μ l/well) on the plate. Dose response curve for Human LRP-5, mFc Tag with the EC50 of 0.63 μ g/ml determined by ELISA.