

# Biotinylated Human MADCAM1 Protein

Cat. No. MCM-HM401B

## Description

<b>Source</b>	Recombinant Biotinylated Human MADCAM1 Protein is expressed from HEK293 with His tag and Avi Tag at the C-Terminus. It contains Gln19-Gln317.
<b>Accession</b>	Q13477-1
<b>Molecular Weight</b>	The protein has a predicted MW of 34.31 kDa. Due to glycosylation, the protein migrates to 50-70 kDa based on Tris-Bis PAGE result.
<b>Endotoxin</b>	Less than 1EU per µg by the LAL method.
<b>Purity</b>	> 95% as determined by Tris-Bis PAGE > 95% as determined by HPLC

## Formulation and Storage


<b>Formulation</b>	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
<b>Reconstitution</b>	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
<b>Storage</b>	-20 to -80°C for 12 months as supplied from date of receipt. -20 to -80°C for 3-6 months in unopened state after reconstitution. 2-8°C for 2-7 days after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

## Background

Mucosal addressin cell adhesion molecule-1 (MAdCAM-1) contributes to the recruitment of donor T cells into the mucosal tissues of the recipient after allogeneic hematopoietic stem cell transplantation (aHSCT). The aim of our study was to determine whether selected single nucleotide polymorphisms (SNPs) of the MADCAM1 gene are associated with development of serious complications after aHSCT.

## Assay Data

### Tris-Bis PAGE




Biotinylated Human MADCAM1 on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.

### SEC-HPLC

# Biotinylated Human MADCAM1 Protein

Cat. No. MCM-HM401B

## Assay Data




The purity of Biotinylated Human MADCAM1 is greater than 95% as determined by SEC-HPLC.

## ELISA Data

### Biotinylated Human MADCAM1, His Tag ELISA

0.1 $\mu$ g Anti-MADCAM1 Antibody, hFc Tag Per Well



Immobilized Anti-MADCAM1 Antibody, hFc Tag at 1 $\mu$ g/ml (100 $\mu$ l/well) on the plate. Dose response curve for Biotinylated Human MADCAM1, His Tag with the EC50 of 7.6ng/ml determined by ELISA.

Log Biotinylated Human MADCAM1, His Tag Conc. ( $\mu$ g/ml)