Mouse HGF R/c-MET Protein

Cat. No. MET-MM101



Description	
Source	Recombinant Mouse HGF R/c-MET Protein is expressed from HEK293 with His tag at the C-terminus.
	It contains Glu25-Ala931.
Accession	P16056
Molecular Weight	The mature form of HGF R is a heterodimer which can be cleaved into α and β chain. The protein has a predicted MW of 32.1 kDa (α chain) and 69.8 kDa (β chain). Due to glycosylation, the protein migrates to 40-50 kDa (α subunit) and 78-98 kDa (β subunit) based on Bis-Tris PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
	> 95% as determined by HPLC
Formulation and Storage	
Formulation	Lyophilized from 0.22 µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.

Background

minimize freeze-thaw cycles.

Storage


c-Met, also called tyrosine-protein kinase Met or hepatocyte growth factor receptor (HGFR), is a protein that in humans is encoded by the MET gene. The protein possesses tyrosine kinase activity. The primary single chain precursor protein is post-translationally cleaved to produce the alpha and beta subunits, which are disulfide linked to form the mature receptor. Following activation by ligand, interacts with the PI3-kinase subunit PIK3R1, PLCG1, SRC, GRB2, STAT3 or the adapter GAB1.

-20 to -80°C for 12 months as supplied from date of receipt.-80°C for 3-6 months after reconstitution.2-8°C for 2-7

days after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please

Assay Data

Bis-Tris PAGE




Mouse HGF R on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

KAGTUS


Assay Data



The purity of Mouse HGF R is greater than 95% as determined by SEC-HPLC.

ELISA Data

Mouse HGF R, His Tag ELISA 0.1µg Mouse HGF R, His Tag Per Well



Immobilized Mouse HGF R, His Tag at $1\mu g/ml$ (100 $\mu l/well$) on the plate. Dose response curve for Human HGF, hFc Tag with the EC50 of 7.6ng/ml determined by ELISA.