Human HLA-A*11:01&B2M&KRAS G12S (VVVGASGVGK) Monomer Protein

Cat. No. MHC-HM442

Description

Source	Recombinant Human HLA-A*11:01&B2M&KRAS G12S (VVVGASGVGK) Monomer Protein is expressed from HEK293 with His tag and Avi tag at the C-Terminus.
	It contains Gly25-Thr305(HLA-A*11:01), Ile21-Met119(B2M) and VVVGASGVGK peptide.
Accession	AAV53343.1(HLA-A*11:01)&P61769(B2M)&VVVGASGVGK
Molecular Weight	The protein has a predicted MW of 50.30 kDa. Due to glycosylation, the protein migrates to 53-63 kDa based on Tris-Bis PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Tris-Bis PAGE
	> 95% as determined by HPLC

Formulation and Storage

Formulation	Lyophilized from 0.22μm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 μg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt.-20 to -80°C for 3-6 months in unopened state after reconstitution.2-8°C for 2-7 days after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

Kirsten rat sarcoma 2 viral oncogene homolog (KRAS) is the most commonly mutated oncogene in human cancer. The developments of many cancers depend on sustained expression and signaling of KRAS, which makes KRAS a high-priority therapeutic target. The virtual screening approach to discover novel KRAS inhibitors and synthetic lethality interactors of KRAS are discussed in detail.

Assay Data

Tris-Bis PAGE



Human HLA-A*11:01&B2M&KRAS G12S (VVVGASGVGK) Monomer on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

For Research Use Only

Human HLA-A*11:01&B2M&KRAS G12S (VVVGASGVGK) Monomer Protein

Cat. No. MHC-HM442

Assay Data



The purity of Human HLA-A*11:01&B2M&KRAS G12S (VVVGASGVGK) Monomer is greater than 95% as determined by SEC-HPLC.

For Research Use Only