## Cynomolgus MICA Protein

Cat. No. MIC-CM10A



Description	
Source	Recombinant Cynomolgus MICA Protein is expressed from HEK293 with His tag at the C-Terminus.
	It contains Glu1-Trp284.
Accession	AAO24115.1
Molecular Weight	The protein has a predicted MW of 33.53 kDa. Due to glycosylation, the protein migrates to 48-58 kDa based on Tris-Bis PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Tris-Bis PAGE
	> 95% as determined by HPLC
Compulation one	1 04

#### Formulation and Storage


romulation and Storage		
Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.	
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 $\mu$ g/ml is recommended. Dissolve the lyophilized protein in distilled water.	
Storage	-20 to -80°C for 12 months as supplied from date of receipt20 to -80°C for 3-6 months in unopened state after reconstitution.2-8°C for 2-7 days after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.	

## **Background**

MICA (MHC class I chain-related gene A) is a transmembrane glycoprotein that functions as a ligand for human NKG2D. A closely related protein, MICB, shares 85% amino acid identity with MICA. These proteins are distantly related to the MHC class I proteins. They possess three extracellular Iglike domains, but they have no capacity to bind peptide or interact with beta 2-microglobulin..

### **Assay Data**

#### Tris-Bis PAGE




Cynomolgus MICA on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC




### **Assay Data**



The purity of Cynomolgus MICA is greater than 95% as determined by SEC-HPLC.

#### **ELISA Data**

# Cynomolgus MICA, His Tag ELISA 0.1µg Cynomolgus MICA, His Tag Per Well



Immobilized Cynomolgus MICA, His Tag at  $1\mu g/ml$  (100 $\mu l/well$ ) on the plate. Dose response curve for Human NKG2D, hFc Tag with the EC50 of 0.14 $\mu g/ml$  determined by ELISA.