

Mouse CD10/MME Protein

Cat. No. MME-MM110

Description

| | |
|-------------------------|---|
| Source | Recombinant Mouse CD10/MME Protein is expressed from HEK293 with His tag at the N-Terminus. It contains Tyr52-Trp750. |
| Accession | Q61391 |
| Molecular Weight | The protein has a predicted MW of 81.1 kDa. Due to glycosylation, the protein migrates to 82-110 kDa based on Bis-Tris PAGE result. |
| Endotoxin | Less than 1EU per μg by the LAL method. |
| Purity | > 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC |

Formulation and Storage


| | |
|-----------------------|---|
| Formulation | Lyophilized from 0.22 μm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization. |
| Reconstitution | Centrifuge the tube before opening. Reconstituting to a concentration more than 100 $\mu\text{g}/\text{ml}$ is recommended. Dissolve the lyophilized protein in distilled water. |
| Storage | -20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles. |

Background

CD10 is an endopeptidase that degrades various bioactive peptides in the extracellular matrix. In addition to enzymatic degradation, it affects multiple intracellular signal transduction pathways. CD10 expression has been extensively studied in human epithelial cancers of numerous organs and sites. CD10 expression pattern depended on the histotypes of thyroid lesions. When possible metastatic tumours and non-epithelial tumours are excluded, high CD10 expression may be useful in determining whether a primary thyroid carcinoma includes an anaplastic component.


Assay Data

Bis-Tris PAGE



Mouse CD10 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC



The purity of Mouse CD10 is greater than 95% as determined by SEC-HPLC.

Mouse CD10/MME Protein

Cat. No. MME-MM110



Assay Data

Bioactivity Data

Measured by its ability to cleave the fluorogenic peptide substrate, Mca-RPPGFSAFK (Dnp) -OH. The specific activity is > 3000 pmoles/min/μg.