

Mouse MMP-9 Protein

Cat. No. MMP-MM109

Description

Source	Recombinant Mouse MMP-9 Protein is expressed from HEK293 with His tag at the C-Terminus. The protein needs to be activated by APMA to have hydrolytic activity. It contains Ala20-Pro730.
Accession	P41245
Molecular Weight	The protein has a predicted MW of 79.64 kDa. Due to glycosylation, the protein migrates to 80-110 kDa based on Tris-Bis PAGE result.
Endotoxin	Less than 1EU per µg by the LAL method.
Purity	> 95% as determined by Tris-Bis PAGE > 95% as determined by HPLC

Formulation and Storage


Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -20 to -80°C for 3-6 months in unopened state after reconstitution. 2-8°C for 2-7 days after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

Matrix metalloproteinase 9 (MMP9) contributes to this process and deficiencies in the MMP9 lead to impaired healing. Inappropriate expression of MMP9 also contributes to impaired re-epithelialization. Previously we demonstrated that FOXO1 was activated in wound healing but to higher levels in diabetic wounds. To address mechanisms of impaired re-epithelialization we examined MMP9 expression in vivo in full thickness dermal scalp wounds created in experimental K14.

Assay Data


Tris-Bis PAGE



Mouse MMP-9 on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

Assay Data



The purity of Mouse MMP-9 is greater than 95% as determined by SEC-HPLC.