

Human Nectin-1/PVRL1/CD111 Protein

Cat. No. NEC-HM201

Description

Source	Recombinant Human Nectin-1/PVRL1/CD111 Protein is expressed from HEK293 with hFc tag at the C-terminus. It contains Gln31-Thr334.
Accession	Q15223-1
Molecular Weight	The protein has a predicted MW of 59.89 kDa. Due to glycosylation, the protein migrates to 70-80 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC

Formulation and Storage


Formulation	Lyophilized from 0.22 μm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 $\mu\text{g}/\text{ml}$ is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3-6 months after reconstitution. 2-8°C for 2-7 days after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

Nectins are a small family of Ca²⁺-independent immunoglobulin (Ig)-like cell adhesion molecules (CAMs) that organize intercellular junctions. They are highly homologous to the human receptor for poliovirus, and as such have been alternately named poliovirus receptor-related proteins. Nectin-1 promotes cell-cell contacts by forming homophilic or heterophilic trans-dimers. Heterophilic interactions have been detected between Nectin-1 and Nectin-3 and between Nectin-1 and Nectin-4. Has some neurite outgrowth-promoting activity.

Assay Data


Bis-Tris PAGE



Human Nectin-1 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

Assay Data




The purity of Human Nectin-1 is greater than 95% as determined by SEC-HPLC.

ELISA Data

Human Nectin-1, hFc Tag ELISA

0.2 μg Human Nectin-3, His Tag Per Well




Immobilized Human Nectin-3, His Tag at 2 μg/ml (100 μl/well) on the plate. Dose response curve for Human Nectin-1, hFc Tag with the EC50 of 14.7 ng/ml determined by ELISA (QC Test).

ELISA Data

Assay Data

Human Nectin-1, hFc Tag ELISA

0.2µg Human Nectin-4, His Tag Per Well



Immobilized Human Nectin-4, His Tag at 2µg/ml (100µl/well) on the plate. Dose response curve for Human Nectin-1, hFc Tag with the EC50 of 0.27µg/ml determined by ELISA.