

Human Nectin-3/CD113 Protein

Cat. No. NEC-HM403

Description

| | |
|-------------------------|--|
| Source | Recombinant Human Nectin-3/CD113 Protein is expressed from HEK293 with His tag and Avi tag at the C-Terminus. It contains Gly58-Asp400. |
| Accession | Q9NQS3 |
| Molecular Weight | The protein has a predicted MW of 40.6 kDa. Due to glycosylation, the protein migrates to 68-82 kDa based on Bis-Tris PAGE result. |
| Endotoxin | Less than 1EU per µg by the LAL method. |
| Purity | > 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC |

Formulation and Storage


| | |
|-----------------------|---|
| Formulation | Lyophilized from 0.22µm filtered solution in 20mM PB, 250mM NaCl (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization. |
| Reconstitution | Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water. |
| Storage | -20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles. |

Background

The Nectin family has at least four members (Nectin-14), all of which show alternate splicing, a transmembrane (TM) region (except for Nectin-1 gamma), and three extracellular Ig-domains. Nectins are highly homologous to the human receptor for poliovirus, and as such, have been alternatively-named poliovirus receptor-related proteins. They do not, however, appear to bind poliovirus. Nectin-3 (also named PRR3, CD113, and PVRL3) is an 83 kDa, type I TM glycoprotein. Its precursor is 549 amino acids (aa) in length. It contains an extended signal sequence of 57 aa, an extracellular domain (ECD) of 347 aa, a transmembrane segment of 21 aa (aa 405-425), and a cytoplasmic region of 124 amino acids.

Assay Data


Bis-Tris PAGE



Human Nectin-3 on Bis-Tris PAGE under reduced conditions. The purity is greater than 95%.

SEC-HPLC

Assay Data




The purity of Human Nectin-3 is greater than 95% as determined by SEC-HPLC.

Assay Data

ELISA Data


Human Nectin-3, His Tag ELISA

0.2µg Human Nectin-3, His Tag Per Well



Immobilized Human Nectin-3, His Tag at 2µg/ml (100µl/well) on the plate. Dose response curve for Human Nectin-1, hFc Tag with the EC50 of 14.7ng/ml determined by ELISA (QC Test).

SPR Data



Human Nectin-3, His Tag immobilized on CM5 Chip can bind Human Nectin-1, His Tag with an affinity constant of 0.204 µM as determined in SPR assay (Biacore T200).