Biotinylated Human Nectin-4 Protein

Cat. No. NEC-HM404B




Description	
Source	Recombinant Biotinylated Human Nectin-4 Protein is expressed from HEK293 with His tag and Avi tag at the C-Terminus.
	It contains Gly32-Val351.
Accession	Q96NY8
Molecular Weight	The protein has a predicted MW of 37.2 kDa. Due to glycosylation, the protein migrates to 46-48 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1 EU per μg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
	> 95% as determined by HPLC
Formulation and Storage	
Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Nectin-4 seems to be involved in cell adhesion through trans-homophilic and -heterophilic interactions, the latter including specifically interactions with NECTIN1. Does not act as receptor for alpha-herpesvirus entry into cells.(Microbial infection) Acts as a receptor for measles virus.

Assay Data

Background

Bis-Tris PAGE




Biotinylated Human Nectin-4 on Bis-Tris PAGE under reduced conditions. The purity is greater than 95%.

SEC-HPLC



Assay Data




The purity of Bioyintlated Human Nectin-4 is greater than 95% as determined by SEC-HPLC.

ELISA Data

Biotinylated Human Nectin-4, His Tag ELISA

 $0.05\mu g$ Biotinylated Human Nectin-4, His Tag Per Well



Immobilized Biotinylated Human Nectin-4, His Tag at 0.5µg/ml (100µl/well) on the streptavidin precoated plate (5µg/ml). Dose response curve for Anti-Nectin-4 Antibody, hFc Tag with the EC50 of 1.9ng/ml determined by ELISA.