

Cynomolgus NGAL/Lipocalin-2 Protein

Cat. No. NGL-CM102

Description	
Source	Recombinant Cynomolgus NGAL/Lipocalin-2 Protein is expressed from HEK293 with His tag at the C-Terminus. It contains Gln21-Gly198.
Accession	XP_005580845.3
Molecular Weight	The protein has a predicted MW of 21.51 kDa. Due to glycosylation, the protein migrates to 25-30 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC

Formulation and Storage


Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 24 months as supplied from date of receipt. -80°C for 3-6 months after reconstitution. 2-8°C for 2-7 days after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

Acute kidney injury (AKI) is one of the most common complications of various serious conditions, and early diagnosis is therefore critical for the treatment of AKI. Recent evidence demonstrates that neutrophil gelatinase-associated lipocalin (NGAL) is closely associated with AKI. Several experimental and clinical studies have shown that the expression of urine and serum NGAL increases significantly in AKI. NGAL shows potential to be a new effective early biochemical marker of AKI. Further studies are needed to confirm the significant advantages of NGAL in the diagnosis of early AKI and its value in clinical applications.

Assay Data


Bis-Tris PAGE



Cynomolgus NGAL on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

Assay Data




The purity of Cynomolgus NGAL is greater than 95% as determined by SEC-HPLC.

ELISA Data

Cynomolgus NGAL, His Tag ELISA

0.05µg Cynomolgus NGAL, His Tag Per Well



Immobilized Cynomolgus NGAL, His Tag at 0.5µg/ml (100µl/Well) on the plate. Dose response curve for Anti-NGAL Antibody, hFc Tag with the EC50 of 3.5ng/ml determined by ELISA (QC Test).