Cynomolgus NGAL/Lipocalin-2 Protein

Cat. No. NGL-CM102



Description	
Source	Recombinant Cynomolgus NGAL/Lipocalin-2 Protein is expressed from HEK293 with His tag at the C-Terminus.
	It contains Gln21-Gly198.
Accession	XP_005580845.3
Molecular Weight	The protein has a predicted MW of 21.51 kDa. Due to glycosylation, the protein migrates to 25-30 kDa based on Tris-Bis PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Tris-Bis PAGE
	> 95% as determined by HPLC

Formulation and Storage


Formulation and Storage	
Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt20 to -80°C for 3-6 months in unopened state after reconstitution.2-8°C for 2-7 days after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

Acute kidney injury (AKI) is one of the most common complications of various serious conditions, and early diagnosis is therefore critical for the treatment of AKI. Recent evidence demonstrates that neutrophil gelatinase-associated lipocalin (NGAL) is closely associated with AKI. Several experimental and clinical studies have shown that the expression of urine and serum NGAL increases significantly in AKI. NGAL shows potential to be a new effective early biochemical marker of AKI. Further studies are needed to confirm the significant advantages of NGAL in the diagnosis of early AKI and its value in clinical applications.

Assay Data

Tris-Bis PAGE




Cynomolgus NGAL on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

KAGTUS


Assay Data



The purity of Cynomolgus NGAL is greater than 95% as determined by SEC-HPLC.

ELISA Data

Cynomolgus NGAL, His Tag ELISA 0.02µg Cynomolgus NGAL, His Tag Per Well



Log Anti-NGAL Antibody, hFc Tag Conc.(µg/ml)

Immobilized Cynomolgus NGAL, His Tag at 0.2µg/ml (100µl/Well) on the plate. Dose response curve for Anti-NGAL Antibody, hFc Tag with the EC50 of 11ng/ml determined by ELISA.