

Cynomolgus NKG2D/CD314 Protein

Cat. No. NKG-CM12D



Description

Source	Recombinant Cynomolgus NKG2D/CD314 Protein is expressed from HEK293 with His tag at the N-Terminus. It contains Ile73-Val216.
Accession	P61252
Molecular Weight	The protein has a predicted MW of 17.48 kDa. Due to glycosylation, the protein migrates to 40-50 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC

Formulation and Storage


Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

NKG2D is a type II transmembrane glycoprotein having an extracellular lectin-like domain. This domain lacks the recognizable calcium-binding sites found in true C-type lectins and binds protein rather than carbohydrate ligands. Human NKG2D is expressed on CD8 alpha beta T cells, gamma δ T cells, NK cells and NKT cells.


Assay Data

Bis-Tris PAGE



Cynomolgus NKG2D on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC




The purity of Cynomolgus NKG2D is greater than 95% as determined by SEC-HPLC.

Assay Data

ELISA Data

Cynomolgus NKG2D, His Tag ELISA

0.05µg Cynomolgus NKG2D, His Tag Per Well



Immobilized Cynomolgus NKG2D, His Tag at 0.5µg/ml (100µl/well) on the plate. Dose response curve for Anti-NKG2D Antibody, hFc Tag with the EC50 of 2.1ng/ml determined by ELISA.