Human NKG2C/CD159c Protein

Cat. No. NKG-HM42C




Description	
Source	Recombinant Human NKG2C/CD159c Protein is expressed from HEK293 with His tag and Avi tag at the N-Terminus.
	It contains Glu98-Leu231
Accession	P26717
Molecular Weight	The protein has a predicted MW of 15.3 kDa. Due to glycosylation, the protein migrates to 40-55 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
Formulation and S	torage
Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.
Background	
	As a first line of defense, natural killer (NK) cells play a crucial role in the fight against infections. The presented study is the first of its kind that ascribes CD56dimCD16 NKG2C-expressing NK cells a crucial role in biasing adaptive immune responses upon influenza vaccination and suggests NKG2C as a potential biomarker in

predicting pandemic influenza vaccine responsiveness.

Assay Data

Bis-Tris PAGE



Human NKG2C/CD159c on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.