Cynomolgus NKp46/NCR1/CD335 Protein

Cat. No. NKP-CM246



Description		
Source	Recombinant Cynomolgus NKp46/NCR1/CD335 Protein is expressed from HEK293 with hFc tag at the C-Terminus.	
	It contains Pro22-Asn254.	
Accession	Q95JB9	
Molecular Weight	The protein has a predicted MW of 53.1 kDa. Due to glycosylation, the protein migrates to 60-70 kDa based on Tris-Bis PAGE result.	
Endotoxin	Less than 1EU per μg by the LAL method.	
Purity	> 95% as determined by Tris-Bis PAGE	
	> 95% as determined by HPLC	
Formulation and Storage		


Formulation	Lyophilized from 0.22 µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt20 to -80°C for 3-6 months in unopened state after reconstitution. 2-8°C for 2-7 days after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

NKp46, along with NKp30 and NKp44, are activating receptors that have been collectively termed the natural cytotoxicity receptors (NCR). These receptors lack significant sequence homology to one another. They are expressed almost exclusively by NK cells and play a major role in triggering some of the key lytic activities of NK cells.

Assay Data

Tris-Bis PAGE




Cynomolgus NKp46 on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

Cat. No. NKP-CM246



Assay Data




The purity of Cynomolgus NKp46 is greater than 95% as determined by SEC-HPLC.



Assay Data

ELISA Data

Cynomolgus NKp46, hFc Tag ELISA 0.05µg Cynomolgus NKp46, hFc Tag Per Well



Log Biotinylated Anti-Nkp46 Antibody, hFc Tag Conc.(µg/ml)

Immobilized Cynomolgus NKp46, hFc Tag at $0.5\mu g/ml$ (100 $\mu l/well$) on the plate. Dose response curve for Biotinylated Anti-Nkp46 Antibody, hFc Tag with the EC50 of 32.5ng/ml determined by ELISA.