

Mouse Noggin Protein

Cat. No. NOG-MM001



Description

Source	Recombinant Mouse Noggin Protein is expressed from HEK293 without tag. It contains Gln28-Cys232.
Accession	P97466
Molecular Weight	The protein has a predicted MW of 23.07 kDa. Due to glycosylation, the protein migrates to 28-33 kDa based on Tris-Bis PAGE result.
Endotoxin	Less than 1EU per µg by the LAL method.
Purity	> 95% as determined by Tris-Bis PAGE > 95% as determined by HPLC

Formulation and Storage


Formulation	Lyophilized from 0.22µm filtered solution in PBS, 100mM L-arginine (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -20 to -80°C for 3-6 months in unopened state after reconstitution. 2-8°C for 2-7 days after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

Noggin is an antagonist of bone morphogenetic proteins (BMP), being indispensable for certain developmental events. Noggin expression positively correlated with EGFR expression in both GC cell line models and The Cancer Genome Atlas human GC cohort. Targeting EGFR and its downstream pathways diminished cell proliferation which was promoted by Noggin. Noggin promotes the proliferation of GC cells by upregulating EGFR and enhancing a vicious circle formed by β catenin, EGFR, ERK and Akt.


Assay Data

Tris-Bis PAGE



Mouse Noggin on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC



The purity of Mouse Noggin is greater than 95% as determined by SEC-HPLC.


Mouse Noggin Protein

Cat. No. NOG-MM001

Assay Data

Cell Based Assay

Recombinant Mouse Noggin Bioactivity



Measured by its ability to inhibit BMP-4-induced alkaline phosphatase production by ATDC5 mouse chondrogenic cells. The ED50 for this effect is 4-80 ng/mL in the presence of 50 ng/mL of recombinant Human BMP4.