Mouse OSCAR Protein

Cat. No. OAR-MM201



Description	
Source	Recombinant Mouse OSCAR Protein is expressed from HEK293 with hFc tag at the C-Terminus.
	It contains Asp19-Asn228.
Accession	Q8VBT3-1
Molecular Weight	The protein has a predicted MW of 49.7 kDa. Due to glycosylation, the protein migrates to 60-70 kDa based on Tris-Bis PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Tris-Bis PAGE
	> 95% as determined by HPLC
Formulation and	Storage
Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before


1 officiation and clorage	
Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt20 to -80°C for 3-6 months in unopened state after reconstitution.2-8°C for 2-7 days after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

Osteoclast-associated receptor (OSCAR) is a co-stimulatory receptor in osteoclastogenesis. Synovial tissues from active rheumatoid arthritis (RA) patients express higher levels of OSCAR compared with osteoarthritic and normal patients. OSCAR and tartrate-resistant acid phosphatase (TRAP) expression levels did not differ between the cartilage pannus junction (CPJ) and non-CPJ regions in active RA.

Assay Data

Tris-Bis PAGE




Mouse OSCAR on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

Cat. No. OAR-MM201



Assay Data



The purity of Mouse OSCAR is greater than 95% as determined by SEC-HPLC.