

Mouse Periostin/OSF-2 Protein

Cat. No. OSF-MM102

Description

Source	Recombinant Mouse Periostin/OSF-2 Protein is expressed from HEK293 with His tag at the C-Terminus.
	It contains Asn24-Gln811.
Accession	Q62009-1
Molecular Weight	The protein has a predicted MW of 88.7 kDa. Due to glycosylation, the protein migrates to 90-95 kDa based on Tris-Bis PAGE result.
Endotoxin	Less than 1EU per µg by the LAL method.
Purity	> 95% as determined by Tris-Bis PAGE
	> 95% as determined by HPLC

Formulation and Storage


Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3-6 months after reconstitution. 2-8°C for 2-7 days after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

Periostin is a matricellular protein that is expressed in several tissues during embryonic development; however, its expression in adults is mostly restricted to collagen-rich connective tissues. Periostin is expressed only briefly during kidney development, but it is not normally detected in the adult kidney. Recent evidence has revealed that periostin is aberrantly expressed in several forms of chronic kidney disease (CKD), and that its expression correlates with the degree of interstitial fibrosis and the decline in renal function.

Assay Data

Tris-Bis PAGE



Mouse Periostin on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.


SEC-HPLC

Mouse Periostin/OSF-2 Protein

Cat. No. OSF-MM102



Assay Data



The purity of Mouse Periostin is greater than 95% as determined by SEC-HPLC.