

Biotinylated Human OSMR Protein

Cat. No. OSM-HM401B

Description

Source	Recombinant Biotinylated Human OSMR Protein is expressed from HEK293 with His tag and Avi tag at the C-terminus. It contains Glu28-Met740.
Accession	Q99650-1
Molecular Weight	The protein has a predicted MW of 84.03 kDa. Due to glycosylation, the protein migrates to 110-140 kDa based on Tris-Bis PAGE result.
Endotoxin	Less than 1EU per µg by the LAL method.
Purity	> 95% as determined by Tris-Bis PAGE > 95% as determined by HPLC

Formulation and Storage


Formulation	Supplied as 0.22 µm filtered solution in PBS (pH 7.4).
Storage	Valid for 12 months from date of receipt when stored at -80°C. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

OSMR is targeted to the mitochondrial matrix via the presequence translocase-associated motor complex components, mtHSP70 and TIM44. OSMR interacts with NADH ubiquinone oxidoreductase 1/2 (NDUFS1/2) of complex I and promotes mitochondrial respiration. Deletion of OSMR impairs spare respiratory capacity, increases reactive oxygen species, and sensitizes BTSCs to IR-induced cell death.


Assay Data

Tris-Bis PAGE



Biotinylated Human OSMR on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC




The purity of Biotinylated Human OSMR is greater than 95% as determined by SEC-HPLC.

Assay Data

ELISA Data

Biotinylated Human OSMR, His Tag ELISA

0.05µg Biotinylated Human OSMR, His Tag Per Well



Immobilized Biotinylated Human OSMR, His Tag at 0.5µg/ml (100µl/well) on the streptavidin precoated plate (5µg/ml). Dose response curve for Anti-OSMR Antibody, hFc Tag with the EC50 of 4.5ng/ml determined by ELISA.