Biotinylated Human uPAR/PLAUR Protein

Cat. No. PAR-HM401B

Description

Source	Recombinant Biotinylated Human uPAR/PLAUR Protein is expressed from HEK293 with His tag and Avi tag at the C-Terminus.
	It contains Leu23-Gly305.
Accession	Q03405-1
Molecular Weight	The protein has a predicted MW of 34.36 kDa. Due to glycosylation, the protein migrates to 50-65 kDa based on Tris-Bis PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Tris-Bis PAGE
	> 95% as determined by HPLC

Formulation and Storage

Formulation	Lyophilized from 0.22μm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 μg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt.-20 to -80°C for 3-6 months in unopened state after reconstitution.2-8°C for 2-7 days after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

The receptor (u-PAR) for urokinase plasminogen activator (u-PA) is a three-domain protein, GPI-anchored to the cell surface, which focuses the enzymatic activity of u-PA, and allows the cell surface activation of plasminogen.Regulation of the activity of u-PA is also mediated by u-PAR.

Assay Data

Tris-Bis PAGE



Biotinylated Human uPAR on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

Assay Data



The purity of Biotinylated Human uPAR is greater than 95% as determined by SEC-HPLC.

ELISA Data



Immobilized Human PLAU, His Tag at 1μg/ml (100μl/well) on the plate. Dose response curve for Biotinylated Human uPAR, His Tag with the EC50 of 13.4ng/ml determined by ELISA.