

Mouse uPAR/PLAUR Domain (2+3) Protein

Cat. No. PAR-MM1D2

Description

| | |
|-------------------------|--|
| Source | Recombinant Mouse uPAR/PLAUR Domain (2+3) Protein is expressed from HEK293 with His tag and Avi tag at the C-terminus. It contains Leu117-Gly298. |
| Accession | P35456-1 |
| Molecular Weight | The protein has a predicted MW of 22.37 kDa. Due to glycosylation, the protein migrates to 40-60 kDa based on Bis-Tris PAGE result. |
| Endotoxin | Less than 1EU per µg by the LAL method. |
| Purity | > 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC |

Formulation and Storage


| | |
|-----------------------|--|
| Formulation | Lyophilized from 0.22 µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization. |
| Reconstitution | Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water. |
| Storage | -20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3-6 months after reconstitution. 2-8°C for 2-7 days after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles. |

Background

The receptor (u-PAR) for urokinase plasminogen activator (u-PA) is a three-domain protein, GPI-anchored to the cell surface, which focuses the enzymatic activity of u-PA, and allows the cell surface activation of plasminogen. Regulation of the activity of u-PA is also mediated by u-PAR.

Assay Data

Bis-Tris PAGE




Mouse uPAR Domain (2+3) on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

Mouse uPAR/PLAUR Domain (2+3) Protein

Cat. No. PAR-MM1D2

Assay Data



The purity of Mouse uPAR Domain (2+3) is greater than 95% as determined by SEC-HPLC.