Mouse uPAR/PLAUR Domain 3 Protein

minimize freeze-thaw cycles.

Cat. No. PAR-MM1D3



Description	
Source	Recombinant Mouse uPAR/PLAUR Domain 3 Protein is expressed from HEK293 with His tag and Avi tag at the C-terminus.
	It contains Phe213-Gly298.
Accession	P35456-1
Molecular Weight	The protein has a predicted MW of 11.83 kDa. Due to glycosylation, the protein migrates to 25-40 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
	> 95% as determined by HPLC
Formulation and	Storage
Formulation	Lyophilized from $0.22~\mu m$ filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
	-20 to -80°C for 12 months as supplied from date of receipt80°C for 3-6 months after reconstitution.2-8°C for 2-7

Background


Storage

The receptor (u-PAR) for urokinase plasminogen activator (u-PA) is a three-domain protein, GPI-anchored to the cell surface, which focuses the enzymatic activity of u-PA, and allows the cell surface activation of plasminogen. Regulation of the activity of u-PA is also mediated by u-PAR.

days after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please

Assay Data

Bis-Tris PAGE



Mouse uPAR Domain 3 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.


SEC-HPLC

Mouse uPAR/PLAUR Domain 3 Protein

Cat. No. PAR-MM1D3

KAGTUS

Assay Data



The purity of Mouse uPAR Domain 3 is greater than 95% as determined by SEC-HPLC.