

Human PD-L1/B7-H1 Protein, Ultra Low Endotoxin



Cat. No. PDL-HM210-UL


Description	
Source	Recombinant Human PD-L1/B7-H1 Protein is expressed from HEK293 with hFc tag at the C-Terminus. It contains Phe19-Arg238.
Accession	Q9NZQ7-1
Molecular Weight	The protein has a predicted MW of 52 kDa. Due to glycosylation, the protein migrates to 68-80 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 0.01 EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC

Formulation and Storage	
Formulation	Lyophilized from 0.22 µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background	
B7-H1, also known as PD-L1 and CD274, is an approximately 65 kDa transmembrane glycoprotein in the B7 family of immune regulatory molecules. PD-L1 has been identified as the ligand for the immunoinhibitory receptor programmed death-1(PD1/PDCD1) and has been demonstrated to play a role in the regulation of immune responses and peripheral tolerance.	

Assay Data


Bis-Tris PAGE



Human PD-L1 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

Assay Data




The purity of Human PD-L1 is greater than 95% as determined by SEC-HPLC.

ELISA Data


Human PD-L1, hFc Tag ELISA

0.2µg Human PD-1, mFc Tag Per Well



Immobilized Human PD-1, mFc Tag at 2 µg/ml (100 µl/well) on the plate. Dose response curve for Human PD-L1, hFc Tag with the EC50 of 0.12 µg/ml determined by ELISA (QC Test).

SPR Data



Human PD-L1, hFc Tag captured on CM5 Chip via Protein A can bind Human PD-1, His Tag with an affinity constant of 2.20 µM as determined in SPR assay (Biacore T200).