Biotinylated Human PD-L2/B7-DC Protein





Description	
Source	Recombinant Biotinylated Human PD-L2/B7-DC Protein is expressed from HEK293 with His tag and Avi tag at the C-Terminus.
	It contains Leu20-Thr220.
Accession	Q9BQ51-1
Molecular Weight	The protein has a predicted MW of 25.3 kDa. Due to glycosylation, the protein migrates to 40-52 kDa based on Tris-Bis PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Tris-Bis PAGE
	> 95% as determined by HPLC
Formulation and	d Storage
E	Lyophilized from 0.22um filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before


Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trenalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 μ g/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt20 to -80°C for 3-6 months in unopened state after reconstitution.2-8°C for 2-7 days after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

PD-1 ligand 2 (PD-L2) as a second ligand for PD-1 and compare the function and expression of PD-L1 and PD-L2. Engagement of PD-1 by PD-L2 dramatically inhibits T cell receptor (TCR)-mediated proliferation and cytokine production by CD4 T cells. At low antigen concentrations, PD-L2-PD-1 interactions inhibit strong B7-CD28 signals.

Assay Data

Tris-Bis PAGE




Biotinylated Human PD-L2 on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC




Assay Data



The purity of Biotinylated Human PD-L2 is greater than 95% as determined by SEC-HPLC.

SPR Data



Human PD-1, hFc Tag captured on CM5 Chip via Protein A can bind Biotinylated Human PD-L2, His-Avi Tag with an affinity constant of 0.344 μ M as determined in SPR assay (Biacore T200).