### Human PD-L1/B7-H1 Protein

Cat. No. PDL-HM610



Description	
Source	Recombinant Human PD-L1/B7-H1 Protein is expressed from HEK293 with Llama Fc tag at the C-terminus.
	It contains Phe19-Arg238.
Accession	Q9NZQ7-1
Molecular Weight	The protein has a predicted MW of 52.50 kDa. Due to glycosylation, the protein migrates to 65-75 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
	> 95% as determined by HPLC

#### Formulation and Storage


Formulation and Storage	
Formulation	Lyophilized from 0.22 µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

B7-H1, also known as PD-L1 and CD274, is an approximately 65 kDa transmembrane glycoprotein in the B7 family of immune regulatory molecules. PD-L1 has been identified as the ligand for the immunoinhibitory receptor programmed death-1(PD1/PDCD1) and has been demonstrated to play a role in the regulation of immune responses and peripheral tolerance.


### **Assay Data**

### **Bis-Tris PAGE**



Human PD-L1 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

## **SEC-HPLC**




The purity of Human PD-L1 is greater than 95% as determined by SEC-HPLC.



### **Assay Data**

**ELISA Data** 

# **Human PD-L1, Llama Fc Tag ELISA** 0.5μg Human PD-L1, Llama Fc Tag Per Well



Immobilized Human PD-L1, Llama Fc Tag at  $5\mu g/ml$  (100 $\mu l/well$ ) on the plate. Dose response curve for Human PD-1, mFc Tag with the EC50 of 0.17 $\mu g/ml$  determined by ELISA.