

Cynomolgus PGF Protein

Cat. No. PGF-CM101

Description

Source	Recombinant Cynomolgus PGF Protein is expressed from HEK293 with His tag at the C-Terminus. It contains Ala21-Arg169.
Accession	A0A2K5VMT8
Molecular Weight	The protein has a predicted MW of 18.16 kDa. Due to glycosylation, the protein migrates to 25-35 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE

Formulation and Storage


Formulation	Lyophilized from 0.22 μm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 $\mu\text{g}/\text{ml}$ is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

Placental growth factor (PGF) is another member of the VEGF family of cytokines with pro-angiogenic and pro-inflammatory effects. Retinal inhibition of PGF in combination with VEGF-A prevents vascular leakage and CNV possibly via modulating their own expression in mononuclear phagocytes. PGF-related, optimized strategies to target inflammation-mediated angiogenesis may help to increase efficacy and reduce non-responders in the treatment of wet AMD patients.

Assay Data


Bis-Tris PAGE



Cynomolgus PGF on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

ELISA Data

Cynomolgus PGF, His Tag ELISA
0.05 μg Cynomolgus PGF, His Tag Per Well



Immobilized Cynomolgus PGF, His Tag at 0.5 $\mu\text{g}/\text{ml}$ (100 $\mu\text{l}/\text{well}$) on the plate. Dose response curve for Anti-PGF Antibody, hFc Tag with the EC₅₀ of 6.0ng/ml determined by ELISA.