

Biotinylated Human PLAU/uPA Protein (active form)

Cat. No. PLA-HM401B

Description

Recombinant Biotinylated Human PLAU/uPA Protein (active form) is expressed from HEK293 with His tag and Avi tag at the C-Terminus.

Source

It contains Ser21-Leu431, which consists of two chains: Long chain A (Ser21-Phe177) and chain B (Ile179-Leu 431). The long chain A is further cleaved to yield a short chain A (Lys156-Phe 177) and N-Terminus fragment (Ser21-Lys155).

Accession

P00749-1

Molecular Weight

The protein has a predicted MW of 49.3 kDa. Due to protein lysis and glycosylation, the protein migrates to 24-26 kDa, 35-38 kDa and 52-60 kDa based on Bis-Tris PAGE result.

Endotoxin

Less than 1EU per ug by the LAL method.

Purity

> 95% as determined by Bis-Tris PAGE

> 95% as determined by HPLC

Formulation and Storage

Formulation

Supplied as 0.22µm filtered solution in PBS (pH 7.4).

Storage

Valid for 12 months from date of receipt when stored at -80°C. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

Plasminogen activator, urokinase (uPA) is a secreted serine protease whose Dysregulation is often accompanied by various cancers. PLAU inhibition could suppress tumor growth. Collectively, PLAU is necessary for tumor progression and can be a diagnostic and prognostic biomarker in HNSCC.


Assay Data

Bis-Tris PAGE



Biotinylated Human PLAU on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC



The purity of Biotinylated Human PLAU is greater than 95% as determined by SEC-HPLC.

Biotinylated Human PLAU/uPA Protein (active form)


Cat. No. PLA-HM401B

Assay Data

ELISA Data

Biotinylated Human PLAU, His Tag ELISA

0.1µg Human uPAR, hFc Tag Per Well



Immobilized Human uPAR, hFc Tag at 1µg/ml (100µl/well) on the plate. Dose response curve for Biotinylated Human PLAU, His Tag with the EC50 of 21.5ng/ml determined by ELISA (QC Test).