

Human PSMA/FOLH1 Protein

Cat. No. PSM-HM210

Description

Source	Recombinant Human PSMA/FOLH1 Protein is expressed from HEK293 with hFc tag at the N-Terminus. It contains Lys44-Ala750.
Accession	Q04609-1
Molecular Weight	The protein has a predicted MW of 106.8 kDa. Due to glycosylation, the protein migrates to 110-120 kDa based on Tris-Bis PAGE result.
Endotoxin	Less than 1EU per µg by the LAL method.
Purity	> 95% as determined by Tris-Bis PAGE > 95% as determined by HPLC

Formulation and Storage


Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -20 to -80°C for 3-6 months in unopened state after reconstitution. 2-8°C for 2-7 days after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

Prostate-specific membrane antigen (PSMA) is an enzyme that in humans is encoded by the FOLH1 (folate hydrolase 1) gene, also known as Glutamate carboxypeptidase II (GCPII). Human PSMA is highly expressed in the prostate, roughly a hundred times greater than in most other tissues. In some prostate cancers, PSMA is the second-most upregulated gene product, with an 8- to 12-fold increase over levels in noncancerous prostate cells.

Assay Data


Tris-Bis PAGE



Human PSMA on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC


Assay Data



The purity of Human PSMA is greater than 95% as determined by SEC-HPLC.

ELISA Data

Human PSMA, hFc Tag ELISA
0.2µg Human PSMA, hFc Tag Per Well



Immobilized Human PSMA, hFc Tag at 2µg/ml (100µl/Well) on the plate. Dose response curve for Biotinylated Anti-PSMA Antibody, hFc Tag with the EC50 of 16.3ng/ml determined by ELISA.