
## **Canine RNF43 Protein**

## Cat. No. RNF-DM134

## ϗͶϲϿ·ႮႽ

| Description             |   |  |  |  |
|-------------------------|---|--|--|--|
| Source                  | Recombinant Canine RNF43 Protein is expressed from HEK293 with His tag at the C-terminus.   |  |  |  |
| Source                  | It contains Gly24-Tyr197.   |  |  |  |
| Accession               | XP_038533015.1  |  |  |  |
| Molecular<br>Weight     | The protein has a predicted MW of 20.68 kDa. Due to glycosylation, the protein migrates to 30-40 kDa based on<br>Bis-Tris PAGE result.  |  |  |  |
| Endotoxin               | Less than 1EU per μg by the LAL method.   |  |  |  |
| Purity                  | > 95% as determined by Bis-Tris PAGE  |  |  |  |
| Funty                   | > 95% as determined by HPLC   |  |  |  |
| Formulation and Storage |   |  |  |  |
| Formulation             | Lyophilized from 0.22 µm filtered solution in PBS (pH 7.4); Normally 8% trehalose is added as protectant before lyophilization.   |  |  |  |
| Reconstitution          | Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.  |  |  |  |
| Storage                 | -20 to -80°C for 12 months as supplied from date of receipt80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.   |  |  |  |
| Background              |   |  |  |  |
|                         | RNF43 (E3 ubiquitin-protein ligase RNF43 or RING-type E3 ubiquitin transferase RNF43) functions as a tumor suppressor, by exerting a predominant negative feedback mechanism in the Wnt/β-catenin signaling pathway. RNF43 inhibits Wnt/beta-catenin signaling by ubiquitinating Frizzled receptor and targeting it to the lysosomal pathway for degradation. |  |  |  |
| Assay Data              |   |  |  |  |

## **Bis-Tris PAGE**



Canine RNF43 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

| Canine RNF43 Protein |                                       |                                       | ͶΛϤϽϓ   |
|----------------------|---------------------------------------|---------------------------------------|---|
| Cat. No.             | RNF-DM134                             |                                       |   |
| Assay Da             | ata                                   |                                       |   |
| mAU  <br>30          |                                       | 9-66.2                                |   |
| 25 -                 |                                       |                                       |   |
| 20 -                 |                                       |                                       |   |
| 15 -                 |                                       |                                       | The purity of Canine RNF43 is greater than 95% as determined by SEC-HPLC. |
| 10 -                 |                                       |                                       |   |
| 5-                   |                                       |                                       |   |
|                      | · · · · · · · · · · · · · · · · · · · | · · · · · · · · · · · · · · · · · · · |   |
| Ō                    | 2 4 6                                 | 8 10 12                               | 14 min  |