

Cat. No. ROR-HM10K

Description


Source	Recombinant Human/Cynomolgus/Rhesus macaque ROR1 (308-395, Kringle Domain) Protein is expressed from HEK293 with His tag at the C-Terminus. It contains Asn308-Asp395.
Accession	Q01973-1(Human) / A0A2K5WTX7(Cynomolgus) / F6RUP2(Rhesus macaque)
Molecular Weight	The protein has a predicted MW of 11.1 kDa. Due to glycosylation, the protein migrates to 17-20 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC

Formulation and Storage

Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background


ROR1 (Receptor tyrosine kinase-like orphan receptor 1), also known as neurotrophic tyrosine kinase receptor-related 1 (NTRKR1), is a member of the ROR family within receptor tyrosine kinases (RTK) superfamily. Two ROR family members (ROR1 and ROR2) have been identified and are characterized by the intracellular tyrosine kinase domains, highly related to those of the Trk-family receptor tyrosine kinases, and by the extracellular Frizzled-like cysteine-rich domains and kringle domains, which are common to receptors of the Wnt family members.

Assay Data**Bis-Tris PAGE**

Human/Cynomolgus/Rhesus macaque ROR1 (308-395, Kringle Domain) on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

Assay Data



The purity of Human/Cynomolgus/Rhesus macaque ROR1 (308-395, Kringle Domain) is greater than 95% as determined by SEC-HPLC.