## **Human SCF Protein**

Cat. No. SCF-HE001



Description	
Source	Recombinant Human SCF Protein is expressed from E.coli without tag.
	It contains Glu26-Ala189.
Accession	P21583-1
Molecular Weight	The protein has a predicted MW of 18.45 kDa same as Tris-Bis PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Tris-Bis PAGE

# Formulation and Storage

Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 μg/ml is recommended. Dissolve the lyophilized protein in distilled water.
	-20 to -80°C for 12 months as supplied from date of receipt80°C for 3-6 months after reconstitution.2-8°C for 2-7

# Background


Storage

Kit-ligand (Kitl) also known as steel factor, stem cell factor and mast cell growth factor plays a crucial role in the development and maintenance of the melanocyte lineage in adult skin. Kitl exerts permanent survival, proliferation and migration functions in Kit receptor-expressing melanocytes.

days after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please

### **Assay Data**

#### Tris-Bis PAGE




Human SCF on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.

## **ELISA Data**

#### Human SCF, No Tag ELISA 0.5μg Human SCF, No Tag Per Well

minimize freeze-thaw cycles.




Immobilized Human SCF at  $5\mu g/ml$  ( $100\mu l/well$ ) on the plate. Dose response curve for Human CD117, His Tag with the EC50 of  $3.42\mu g/ml$  determined by ELISA.



# **Assay Data**

### **Cell Based Assay**

# **Recombinant Human SCF Bioactivity**



Measured in a cell proliferation assay using Mo7e cells. The ED50 for this effect is 1-8 ng/mL (QC Test).