Biotinylated Human Semaphorin 4D/SEMA4D/CD100 Protein






Cat. NO. SLIVI-I II	MITTO
Description	
Source	Recombinant Biotinylated Human Semaphorin 4D/SEMA4D/CD100 Protein is expressed from HEK293 with His tag and Avi tag at the C-Terminus.
	It contains Met22-Arg734.
Accession	Q92854-1
Molecular Weight	The protein has a predicted MW of 82.1 kDa. Due to glycosylation, the protein migrates to 115-140 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
	> 95% as determined by HPLC
Formulation and S	Storage
Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.
Background	
	Semaphorin 4D (Sema4D) is a multifunctional protein widely expressed in an organism that plays an important

role in the control of many physiological and pathological processes, including immunoregulation, neurogenesis,


angiogenesis, and tumor progression. **Assay Data**

Bis-Tris PAGE



Biotinylated Human Semaphorin 4D on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC



The purity of Biotinylated Human Semaphorin 4D is greater than 95% as determined by SEC-HPLC.

Biotinylated Human Semaphorin 4D/SEMA4D/CD100 Protein

Cat. No. SEM-HM44DB




Assay Data

ELISA Data

Biotinylated Human Semaphorin 4D, His Tag ELISA

0.1μg Biotinylated Human Semaphorin 4D, His Tag Per Well



Immobilized Biotinylated Human Semaphorin 4D, His Tag at 1 μ g/ml (100 μ l/well) on the streptavidin precoated plate (5 μ g/ml). Dose response curve for Anti-Semaphorin 4D Antibody , hFc Tag with the EC50 of 6.9ng/ml determined by ELISA.