Cynomolgus SG3/Secretogranin 3 Protein, Ultra Low Endotoxin

Cat. No. SGS-CM101-UL

Description

Source	Recombinant Cynomolgus SG3/Secretogranin 3 Protein is expressed from HEK293 with His tag at the C-Terminus.
	It contains Phe20-Leu468.
Accession	I7G9U8
Molecular Weight	The protein has a predicted MW of 52.14 kDa. Due to glycosylation, the protein migrates to 58-70 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 0.01 EU per μg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
	> 95% as determined by HPLC

Formulation and Storage

Formulation	Lyophilized from 0.22μm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 24 months as supplied from date of receipt.-80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

Secretogranin III (Scg3) is a member of the granin protein family that regulates the biogenesis of secretory granules. Scg3 was recently discovered as an angiogenic factor, expanding its functional role to extrinsic regulation. Unlike many other known angiogenic factors, the pro-angiogenic actions of Scg3 are restricted to pathological conditions. Among thousands of quantified endothelial ligands, Scg3 has the highest binding activity ratio to diabetic vs.

Assay Data

Bis-Tris PAGE



Cynomolgus SG3 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

Cynomolgus SG3/Secretogranin 3 Protein, Ultra Low Endotoxin

Cat. No. SGS-CM101-UL

Assay Data



The purity of Cynomolgus SG3 is greater than 95% as determined by SEC-HPLC.

ELISA Data



Immobilized Cynomolgus SG3, His Tag at 1μg/ml (100μl/well) on the plate. Dose response curve for Anti-SG3 Antibody, hFc Tag with the EC50 of 3.4ng/ml determined by ELISA.