

Human CD209/DC-SIGN Protein, Ultra Low Endotoxin



Cat. No. SIG-HM101-UL

Description

Source	Recombinant Human CD209/DC-SIGN Protein is expressed from HEK293 with His tag at the N-Terminus. It contains Gln59-Ala404.
Accession	Q9NNX6-1
Molecular Weight	The protein has a predicted MW of 40.5 kDa. Due to glycosylation, the protein migrates to 52-60 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 0.01 EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC

Formulation and Storage


Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

C-type lectin CD209/DC-SIGN and CD209L/L-SIGN proteins are distinct cell adhesion and pathogen recognition receptors that mediate cellular interactions and recognize a wide range of pathogens, including viruses such as SARS, SARS-CoV-2, bacteria, fungi and parasites. Pathogens exploit CD209 family proteins to promote infection and evade the immune recognition system.

Assay Data


Bis-Tris PAGE



Human CD209 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.


SEC-HPLC

Assay Data



The purity of Human CD209 is greater than 95% as determined by SEC-HPLC.

SPR Data



Human BTN2A1, His Tag immobilized on CM5 Chip can bind Human CD209, His Tag with an affinity constant of 0.12 μ M as determined in SPR assay (Biacore T200).