

Human Siglec-2/CD22 Domain 6&7 Protein, Ultra Low Endotoxin

Cat. No. SIG-HM123-UL

Description

Source	Recombinant Human Siglec-2/CD22 Domain 6&7 Protein is expressed from HEK293 with His tag at the C-terminus. It contains Pro505-Arg687.
Accession	P20273-1
Molecular Weight	The protein has a predicted MW of 22.22 kDa. Due to glycosylation, the protein migrates to 35-40 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 0.01 EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC

Formulation and Storage


Formulation	Lyophilized from 0.22 µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

CD22, or cluster of differentiation-22, is a molecule belonging to the SIGLEC family of lectins. It is found on the surface of mature B cells and to a lesser extent on some immature B cells. CD22 a member of the immunoglobulin superfamily. CD22 functions as an inhibitory receptor for B cell receptor (BCR) signaling. It is also involved in the B cell trafficking to Peyer's patches in mice.

Assay Data


Bis-Tris PAGE



Human Siglec-2 Domain 6&7 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

Assay Data



The purity of Human Siglec-2 Domain 6&7 is greater than 95% as determined by SEC-HPLC.