
# Biotinylated Human Siglec-4a/MAG Protein





Description	
Source	Recombinant Biotinylated Human Siglec-4a/MAG Protein is expressed from HEK293 with His tag and Avi tag at the C-Terminus.
	It contains Gly20-Pro516.
Accession	P20916
Molecular Weight	The protein has a predicted MW of 57.6 kDa. Due to glycosylation, the protein migrates to 78-85 kDa based on Tris-Bis PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Tris-Bis PAGE
	> 95% as determined by HPLC
Formulation and	Storage
Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before


Formulation	Lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt20 to -80°C for 3-6 months in unopened state after reconstitution.2-8°C for 2-7 days after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

**Background** 

Siglec-4a, also known as Myelin-Associated Glycoprotein (MAG), is a type I transmembrane glycoprotein belonging to the Siglec family, a subgroup of the Ig superfamily. Adhesion molecule that mediates interactions between myelinating cells and neurons by binding to neuronal sialic acid-containing gangliosides and to the glycoproteins RTN4R and RTN4RL2 (By similarity).

### **Assay Data**

### Tris-Bis PAGE




Biotinylated Human Siglec-4a on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.

**SEC-HPLC** 

# KAGTUS

### **Assay Data**




The purity of Biotinylated Human Siglec-4a is greater than 95% as determined by SEC-HPLC.

#### **ELISA Data**

## Biotinylated Human Siglec-4a, His Tag ELISA

0.1μg Anti-Siglec-4a Antibody, hFc Tag Per Well



 $\textbf{Log Biotinylated Human Siglec-4a, His Tag Conc.} (\mu g/ml)$ 

Immobilized Anti-Siglec-4a Antibody, hFc Tag at  $1\mu g/ml$  (100 $\mu l/well$ ) on the plate. Dose response curve for Biotinylated Human Siglec-4a, His Tag with the EC50 of 24.4ng/ml determined by ELISA.